

Large-scale collective motion of RFGC galaxies

S. L. Parnovsky • A. S. Parnowski

© Springer-Verlag ●●●●

Abstract We processed the data about radial velocities and HI linewidths for 1678 flat edge-on spirals from the Revised Flat Galaxy Catalogue. We obtained the parameters of the multipole components of large-scale velocity field of collective non-Hubble galaxy motion as well as the parameters of the generalized Tully-Fisher relationship in the “HI line width – linear diameter” version. All the calculations were performed independently in the framework of three models, where the multipole decomposition of the galaxy velocity field was limited to a dipole, quadrupole and octopole terms respectively. We showed that both the quadrupole and the octopole components are statistically significant.

On the basis of the compiled list of peculiar velocities of 1623 galaxies we obtained the estimations of cosmological parameters Ω_m and σ_8 . This estimation is obtained in both graphical form and as a constraint of the value $S_8 = (\Omega_m/0.3)^{0.35}\sigma_8 = 0.91 \pm 0.05$.

Keywords galaxies; large-scale structure; collective motion of galaxies; cosmological parameters

1 Introduction

The distribution of matter, including dark matter, in the Universe is inhomogeneous on the scales of about

100 *Mpc*. This is manifested, for example, in the existence of superclusters and voids. A galaxy, besides the cosmological expansion, is also attracted to the regions with greater density. As a result, the galaxies are involved in a non-Hubble large-scale collective motion on the background of Hubble expansion. In a more sophisticated way this can be considered as a development of initial density and velocity fluctuations in the early Universe due to the gravitational instability.

Investigation of such a motion is important at least for two reasons. First of all, it allows to plot the distribution of matter, including dark matter, in the surrounding region of the Universe, which is the main goal of cosmography, and to compare this distribution with the distribution of luminous matter. The second reason is that the parameters of this motion are linked with certain cosmological parameters, so we can obtain new independent estimations of these parameters. Of course, the accuracy of such estimation will not be very high, but the agreement of different estimations of cosmological parameters can support the correctness of the standard cosmological model.

In 1989 Karachentsev proposed to use flat edge-on spiral galaxies as a tool for studying their large-scale collective motion. They are good in this role for the following reasons:

1. The linear diameter is strongly correlated with the HI linewidth for thin bulgeless galaxies. This allows to determine distances without photometric data.
2. Flat galaxies can be easily identified by their axes ratio.
3. Flat galaxies have a nearly 100 per cent HI detection rate.
4. Galaxies in clusters are usually not flat due to interaction with neighbours. This means that flat galaxies avoid clustering and do not interact with the intergalactic gas in clusters.

S. L. Parnovsky

Astronomical Observatory of Taras Shevchenko Kyiv National University
prov. Observatorny, 3, 04053, Kyiv, Ukraine
tel: +380444860021, fax: +380444862191
e-mail: par@observ.univ.kiev.ua

A. S. Parnowski

Space Research Institute of NASU and NSAU
prosp. Akad. Glushkova, 40, korp. 4/1, 03680 MSP, Kyiv-187, Ukraine
tel: +380933264229, fax: +380445264124
e-mail: parnowski@gmail.com

5. Peculiar velocities of isolated flat galaxies are not perturbed by neighbours.

To implement this idea the Flat Galaxy Catalogue (FGC, Karachentsev et al. 1993) was created. It contained data about $N = 4455$ galaxies, which satisfied the conditions $a_b/b_b \geq 7$ and $a_b > 0.6'$. Here a_b and b_b are the major and minor axial diameters directly measured on POSS-I and ESO/SERC plates. In accordance with the original photographic material, the Catalogue consists of two parts: FGC ($N = 2573$) and its southern extension, FGCE ($N = 1882$). The first part is based on the POSS-I and covers the sky region with declinations between -20 deg and $+90$ deg. The second one is based on the ESO/SERC and covers the rest of the sky area up to $\delta = -90$ deg.

After thorough studies of the catalogue's properties, both these parts were joined. The angular diameters from the FGCE were converted to the POSS-I system of the FGC, which appeared to be close to the a_{25} system. This system, where galaxy size is taken at $B = 25 \text{ mag}/\square''$ isophotal level, was used, in particular, by de Vaucouleurs et al. (1991). Some FGCE galaxies, which did not satisfy the condition $a > 0.6'$ after conversion, were removed from the resulting Revised Flat Galaxy Catalogue (RFGC, Karachentsev et al. 1999). It contained data about $N = 4236$ galaxies including the information on the following parameters: Right Ascension and Declination for the epochs J2000.0 and B1950.0, galactic longitude and latitude, major and minor blue and red diameters in arcminutes in the POSS-I diameter system, morphological type of the spiral galaxies according to the Hubble classification, index of the mean surface brightness (I – high, IV – very low) and some other parameters, which are not used in this article. More detailed description of the catalogue can be found in the paper (Karachentsev et al. 1999).

The original goal of this catalogue was to estimate the distance to galaxies according to the Tully-Fisher relationship in the “HI line width – linear diameter” version without using their redshifts. The difference between the velocity V derived from the redshift and the Hubble velocity $R = Hr$ corresponding to the distance r estimated by Tully-Fisher relationship is called a peculiar velocity $V_{pec} = cz - Hr$. We can use such a simple form of the Hubble's law because our sample has $z < 0.1$.

There are some important things to take into account about peculiar velocities. The redshift includes not only the velocity of the galaxy, but also the velocities of our Galaxy, Solar System and the Earth. Thus, to eliminate these factors, all velocities were reduced to the frame, where CMB is isotropic. Naturally, the redshift gives us only the radial component of the velocity and the tangential components cannot be measured.

Additionally, Tully-Fisher relationship is statistical and thus has a certain error. Thus, we can only estimate the peculiar velocity for each galaxy, sometimes with a significant error.

However, we believe that there is a large-scale velocity field. We consider the individual galaxies as test particles in the velocity field of large-scale collective motion. Thus, having data about the peculiar velocities of a large number of galaxies, we can restore the radial component of the velocity field. Using some additional assumptions, like the potentiality of the flux, it is possible to restore the 3D velocity field. For this reason we need ample samples of peculiar velocities, preferably uniformly covering the celestial sphere.

These conditions are satisfied by the RFGC catalogue, which covers the whole celestial sphere in both hemispheres with the natural exclusion of the Milky Way region. However, not all of the RFGC galaxies have data required to estimate the distance to them using the Tully-Fisher relationship. Nevertheless, the sample of galaxies having such data is also quite uniform, as shown on Fig. 1.

Naturally, RFGC is not the only sample that can be used for this purpose. Possible alternatives include SBF (Tonry et al. 1997, 2000), ENEAR (da Costa et al. 2000a), SFI (Haynes et al. 1999a,b), SFI++ (Springob et al. 2007), EFAR (Wegner et al. 1996; Colless et al. 2001), SMAC (Hudson et al. 1999, 2004), 2MFGC (Mitronova et al. 2003). However, in this article we use only the RFGC catalogue.

To apply the Tully-Fisher relationship the data from the catalogue is not sufficient; we also need to know the HI linewidth W (in this article we take it at 50 per cent level), or the gas rotation velocity V_{rot} obtained from optical observations. Additionally, we need redshift data. The number of galaxies with such data increases constantly.

Since FGC and RFGC were assembled, a number of articles were published dealing with collective motions of RFGC galaxies. Very preliminary results were published by Karachentsev et al. (1995). The parameters of the bulk motion were calculated by Karachentsev et al. (2000b). In the paper (Parnovsky et al. 2001) not only new data were added, but also the models featuring the quadrupole and octopole components of the velocity field were proposed. Also, the generalized Tully-Fisher relationship for RFGC galaxies was finalized. It included data not only about HI linewidth and angular diameter in red and blue bands, but also about the morphological type of the galaxy and its surface brightness index. By that time the authors had information about radial velocities and HI 21 cm line widths or V_{rot} for 1327 RFGC




Fig. 1 Distribution of galaxies over the celestial sphere in galactic coordinates

galaxies from the different sources listed in the paper (Karachentsev et al. 2000b). From this number, 1271 galaxies were included in the sample, and the rest of them were considered to be outliers. As a result, the first list of peculiar velocities of RFGC galaxies was prepared by Karachentsev et al. (2000a). Four years later, the number of galaxies with available data increased and reached 1561 (Parnovsky & Tugay 2004); 1493 of them entered the sample. A new version of the list of peculiar velocities was prepared by Parnovsky & Tugay (2005). This list was the basis for solving the two above-mentioned problems, namely mapping the matter density and estimation of cosmological parameters. The distribution of matter density was obtained in the paper (Sharov & Parnovsky 2006) up to $75h^{-1} \text{ Mpc}$ in the supergalactical plane and 8 more planes. In the same article, the excess masses of attractors in this region were estimated, and the estimation of β parameter was obtained. The estimations of the cosmological parameters Ω_m and σ_8 were obtained in the paper (Parnovsky et al. 2006). The model was expanded taking into account the effects of general theory of relativity in the paper (Parnovsky & Gaydamaka 2004).

An important result, obtained in the paper (Parnovsky et al. 2001), and confirmed by Parnovsky & Tugay (2004), was the statistical significance of quadrupole and octopole components (both at more than 99 per cent confidence level). This result relied on the F-test, which assumes normal distribu-

tion of errors (see Hudson 1964). However, even the normally distributed errors of angular diameters and HI linewidths and deviations from Tully-Fisher relationship lead to non-Gaussian distribution of peculiar velocities. Thus, the statistical significance of these terms required additional consideration. In the article (Parnovsky & Parnowski 2008) it was shown using Monte-Carlo simulations that the statistical significance of these components appeared to be less than perceived from the F-test, but still large enough to be considered: the quadrupole was statistically significant at 96 per cent confidence level and the octopole – at 90.93 per cent confidence level.

In the four years that passed since the last list of peculiar velocities of RFGC galaxies was compiled, not only new data have appeared, but also some old data were remeasured. This led to a necessity of repeating all the steps required to obtain the new list. This means that not only new data have to be added, but the process of sample selection must be repeated from the very beginning. As it will be shown further, some results appeared to be notably different from the ones obtained with the previous sample.

In this paper we present the parameters of the large-scale collective velocity field in the framework of the multipole model as well as the estimations of some cosmological parameters. The distribution of matter density will be discussed in a separate article.

2 Multipole models of collective motion of galaxies

In the paper (Parnovsky et al. 2001) the velocity field components in the CMB reference frame were expanded in terms of galaxy's radial vector $\vec{r}$. The first three terms are

$$V_i = D_i + A_{ik}r_k + O'_{ikl}r_kr_l. \quad (1)$$

In our notation we use the Einstein rule: summation by all the repeating indices. Hence, for the radial velocity V_{3K} in the CMB reference frame we get

$$V_{3K} = D_i n_i + r A_{ik} n_i n_k + r^2 O'_{ikl} n_i n_k n_l, \quad (2)$$

where $\vec{n} = \vec{r}/r$ and $r = |\vec{r}|$. Let us decompose the tensor $\mathbf{A}$ in two parts: $A_{ik} = H\delta_{ik} + Q'_{ik}$. Here H is a trace, corresponding to the Hubble constant, δ_{ik} is the Kronecker delta, and $\mathbf{Q}'$ is a traceless tensor. Next we switch from distance r to the corresponding Hubble velocity $R = Hr$.

$$V_{3K} = R + D_i n_i + RQ_{ik} n_i n_k + R^2 O_{ikl} n_i n_k n_l, \quad (3)$$

where $\mathbf{Q} = H^{-1}\mathbf{Q}'$ and $\mathbf{O} = H^{-2}\mathbf{O}'$.

This decomposition was used to obtain some models of dependence of galaxy's radial velocity V . In the simplest D-model (Hubble law + dipole) we have

$$V_{3K} = R + V^{dip} + \delta V, \quad V^{dip} = D_i n_i, \quad (4)$$

where $\vec{D}$ is a velocity of homogeneous bulk motion, δV is a random deviation and $\vec{n}$ is a unit vector towards galaxy. After the addition of quadrupole terms we obtain a DQ-model

$$V_{3K} = R + V^{dip} + V^{qua} + \delta V, \quad V^{qua} = RQ_{ik} n_i n_k \quad (5)$$

with symmetrical traceless tensor $\mathbf{Q}$ describing quadrupole components of the velocity field. It characterises the relative deviation of an effective "Hubble constant" in a given direction from the mean value. More detailed discussion will be given further. The DQO-model includes the octopole component of velocity field described by a symmetrical tensor $\mathbf{O}$ of rank 3:

$$V_{3K} = R + V^{dip} + V^{qua} + V^{oct} + \delta V, \quad V^{oct} = R^2 O_{ikl} n_i n_k n_l. \quad (6)$$

In some cases it makes sense to use another way of describing the octopole component in the DQO-model. For this purpose the tensor $\mathbf{O}$ can be considered a sum of a traceless tensor $\hat{\mathbf{O}}$, and a trace, characterized by the vector $\vec{P}$

$$O_{ijk} = \hat{O}_{ijk} + P_i \delta_{jk}, \quad P_i = \frac{3}{5} O_{ijk} \delta_{jk}. \quad (7)$$

Here the indices in parantheses are symmetrized. Thus an alternative form of the DQO-model is

$$V_{3K} = R + (D_i + R^2 P_i) n_i + RQ_{ik} n_i n_k + R^2 \hat{O}_{ikl} n_i n_k n_l + \delta V. \quad (8)$$

We will use Cartesian components of the vector $\vec{n}$ in the galactic coordinates:

$$\begin{aligned} n_1 &= n_z = \sin b, \quad n_2 = n_x = \cos l \cos b, \\ n_3 &= n_y = \sin l \cos b. \end{aligned} \quad (9)$$

These three models were used for processing data on RFGC galaxies. To estimate the distances to galaxies a generalized Tully-Fisher relationship was used in the 'linear diameter – HI line width' version. It has a form (Parnovsky et al. 2001)

$$R = (C_1 + C_2 B + C_3 B T + C_4 U) \frac{W}{a} + C_5 \left(\frac{W}{a}\right)^2 + C_6 \frac{1}{a}, \quad (10)$$

where W is a corrected HI line width in km s^{-1} measured at 50 per cent of the maximum, a is a corrected major galaxies' angular diameter in arcminutes on red POSS and ESO/SERC reproductions, U is a ratio of major galaxies' angular diameters on red and blue reproductions, T is a morphological type indicator ($T = I_t - 5.35$, where I_t is a Hubble type; $I_t = 5$ corresponds to type Sc), and B is a surface brightness indicator ($B = I_{SB} - 2$, where I_{SB} is a surface brightness index from RFGC; brightness decreases from I to IV). Details of corrections one can find in the papers (Parnovsky et al. 2001; Parnovsky & Tugay 2004). The reasons for choosing this form of Tully-Fisher relationship for RFGC galaxies are given ibid. We only note that the statistical significance of each term in eq. (10) is greater than 99 per cent.

Note that eq. (10) differs from the classical Tully-Fisher relationship, which has the form

$$R = C \frac{W^\alpha}{a}. \quad (11)$$

The corresponding analysis was performed in the paper (Parnovsky & Gaydamaka 2004) for the relativistic model of motion. It was shown that the power α for the classical form of Tully-Fisher relationship (11) somewhat differs from 1. However, after including the corrections for the surface brightness and the morphological type of the galaxy, the value of α appeared to be close enough to 1 to use the form given by (10). If we repeat this analysis for the new data, the minimal variance is reached at $\alpha \sim 1.16$. At the same time the standard deviation σ for the form (11) is 5.7 per cent more than for the form (10) depending on the depth

of the subsample and model used. If we combine the forms (10) and (11) into

$$R = (C_1 + C_2 B + C_3 B T + C_4 U) \frac{W^\alpha}{a} + C_5 \left(\frac{W^\alpha}{a} \right)^2 + C_6 \frac{1}{a}, \quad (12)$$

we obtain the minimal variance at $\alpha \sim 1.05$, which is very close to 1. This correction does not reduce σ due to the decreased number of degrees of freedom. This makes the introduction of α senseless. For this reason we will use only the form (10). This is additionally justified by the fact that all the coefficients enter this equation linearly and thus we can use the linear regression analysis.

The D-model has 9 parameters (6 coefficients C and 3 components of vector $\vec{D}$), DQ-model has 14 parameters (5 components of tensor $\mathbf{Q}$ are added), and DQO-model is described by 24 coefficients. The values and errors of the coefficients were calculated by the least square method for different subsamples with distances limitation to make the sample more homogeneous in the depth (Parnovsky et al. 2001). We used the same weights for all datapoints. Since the quadrupole and octopole terms explicitly depend on R , an iteration procedure was used. Note that the coefficients of the generalised Tully-Fisher relationship and components of the velocity model were fitted simultaneously. The iterations converge rather quickly.

3 Description of data

Directly from the RFGC catalogue we take the angular diameters on blue and red reproductions, morphological type of a galaxy and its surface brightness index. The angular diameters are corrected for intrinsic absorption. From other sources we take the radial velocity in the CMB frame and the HI line width or the gas rotation velocity V_{rot} obtained from optical observations. The line widths are corrected for cosmological expansion and for turbulence. The data on HI line widths for RFGC galaxies in the HyperLeda (Paturel et al. 2003a) catalogue are based on 149 sources, which have different reliability. The main sources used in previous versions of our samples are listed in the papers (Parnovsky et al. 2001; Parnovsky & Tugay 2004; Karachentsev et al. 2000a,b; Parnovsky & Tugay 2005). Since that time new large volumes of data on radial velocities and HI linewidths of galaxies were published (e.g., ?). For this reason we rechecked the original data on HI line widths of galaxies in previous samples and changed some of them. For each galaxy we used only the data of original measurements and did not average the data of different papers.

In the cases when data from reliable sources differed significantly, we used less reliable sources to choose between them. We also checked the deviation of the distance to galaxy from the Tully-Fisher relationship in the D-model.

After revising data from the previous samples, we added new data measured during the last 5 years. We took HI linewidths from the original articles (Koribalski et al. 2004; Theureau et al. 2005; Doyle et al. 2005; Springob et al. 2005). In the case when W was unavailable we converted V_{rot} from the HyperLeda catalogue (Paturel et al. 2003b) to W using the relationship $W = 2.016 * V_{rot} + 20.6$, obtained by processing data for galaxies, which have both records. Note that there are rare cases of significant deviations from this relationship.

In the process of data preparation we constantly checked the deviations of the radial velocities from the velocities calculated in the D-model. Some data were obvious outliers. Such galaxies were rejected from the sample. As a result, we had all the necessary data for 1678 galaxies, including 55 outliers. Thus the sample contained 1623 galaxies, including 204 new and 1419 old ones. It is noteworthy that some of the old data changed as well. The distribution of these galaxies over the celestial sphere is shown on Fig. 1. The mean radial velocity of old galaxies is 6000 km s^{-1} , and of new ones it is 5600 km s^{-1} . The mode in both cases is about 4800 km s^{-1} . The sample has 90% completeness up to about 6000 km s^{-1} .

The farthest galaxies in the sample have radial velocities above 20000 km s^{-1} . However, at such scales the sample is inhomogeneous and incomplete. To deal with more homogeneous samples, we consider subsamples with distance limitation. They include all the galaxies satisfying the condition $R < R_{max}$ in the D-model and are used for all the three models. For this purpose we used the following iterative procedure. First, we calculate the distances to the galaxies using seed coefficients from the whole sample or another subsample with close R_{max} . Then we select the galaxies with $R < R_{max}$ and recalculate the coefficients using only their data, once again calculate the distances and so on. In most cases these iterations converge, but in some cases, more often for small R_{max} , we get a limit cycle. In this case we select only the galaxies whose distances are below R_{max} for all sets of coefficients in this limit cycle.

We used different subsamples with R_{max} ranging from 3000 to 12000 km s^{-1} . In this paper we mostly present information for the subsamples with R_{max} equal to 8000 km s^{-1} ($r = 80h^{-1} \text{ Mpc}$) and 10000 km s^{-1} ($r = 100h^{-1} \text{ Mpc}$).




Fig. 2 Dependence of the dipole component of the velocity on the sample depth for 3 models. Triangles correspond to the D-model, squares – to the DQ-model, and circles – to the DQO-model

4 Multipole structure of collective motion

The parameters of a generalized Tully-Fisher relationship for all three models and two R_{max} values are given in Table 1 together with their statistical significance levels according to F-test. The F values should be compared to 3.8, 6.6, 7.9, 10.8 and 12.1, which correspond to 95, 99, 99.5, 99.9 and 99.95 per cent confidence levels respectively. Most of them are statistically significant at 99.95 per cent confidence level with the exception of C_3 for $R_{max} = 10000 \text{ km s}^{-1}$. Comparing the coefficients to that of Parnovsky & Tugay (2004) one can see that C_1 , C_4 , C_5 and C_6 remain the same within the margin of error, and C_2 and especially C_3 are reduced, which means lesser influence of the morphological type of galaxies. This also leads to some decrease of statistical significance, which is still more than 95 per cent.

The inclusion of the octopole component for subsamples with R_{max} about 8000 km s^{-1} leads to a drastic decrease of the dipole component up to 1σ level, as Parnovsky et al. (2001) have shown. This effect appeared to be even more evident in the paper (Parnovsky & Tugay 2004). Such a strong dipole-octopole coupling can be explained by the same symmetry of these components with respect to the inversion of space. For the new sample this effect appeared to be weaker as the dipole component decreases only up to 2σ level (see Fig. 2). Naturally this effect is caused by incompleteness and asymmetry of the sam-




Fig. 3 Bulk motion apices in galactic coordinates for $R_{max} = 10000 \text{ km s}^{-1}$. Crosses mark the apices of the bulk motion in the D-model surrounded by 1σ , 2σ and 3σ confidence areas. The square marks the apex position of the galaxies from the previous sample with corrected data. The diamond marks the apex position in the DQO-model. Solid boundaries correspond to new results, and dashed ones – to the results of Parnovsky & Tugay (2004). Numbers denote the results of other authors: 1 – (Lynden-Bell et al. 1988), 2 – (Hudson et al. 1995), 3 – (Lauer & Postman 1994), 4 – (Parnovsky et al. 2001), 5 – (Dekel et al. 1999), 6 – (da Costa et al. 2000b), 7 – (Hudson et al. 2004), 8 – (Dale et al. 1999), 9 – (Kudrya et al. 2003), 10 – (Watkins et al. 2009)

ple, which lead to dipole and octopole components being non-orthogonal.

This effect is important because D and DQO models yield the values of the bulk motion velocity differing by a factor of 2. Thus the dependence of the bulk motion velocity on the sample depth will be totally different for D and DQO models as one can see from Fig. 2. One should take this into account when analysing the convergence scale.

In comparison with the papers (Parnovsky et al. 2001) and (Parnovsky & Tugay 2004), the apex direction of the dipole component has changed. In Table 2 we present its parameters for 2 subsamples together with the standard deviation σ for D-model. Note that σ includes both the intrinsic scatter in the generalised Tully-Fisher relationship and the stochastic component of the velocities, resulting in high values.

The comparison of the results from Table 2 with those obtained earlier (Parnovsky et al. 2001; Parnovsky & Tugay 2004) shows that the norm of the velocity hasn't changed significantly, but the apex direction has. Let us discuss this question in detail. On Fig. 3 we present the Mollweide projection upon the celestial sphere of the ellipsoids corresponding to 1σ , 2σ and 3σ confidence levels for the apex direction in the framework of D-model for $R_{max} = 10000 \text{ km s}^{-1}$. Solid boundaries correspond to new results, and dashed ones – to the results of Parnovsky & Tugay (2004). One can see that the apex direction has changed at slightly more than 2σ . Besides,

Table 1 Parameters of the generalized Tully-Fisher relationship (10)

i	D-model		DQ-model		DQO-model	
	C_i	F_i	c_i	F_i	c_i	F_i
$R_{max} = 8000$ km/s (1240 galaxies)						
1	15.8 ± 1.6	103	17.2 ± 1.6	122	17.4 ± 1.6	126
2	1.84 ± 0.20	80.5	1.96 ± 0.21	90.0	1.95 ± 0.21	88.4
3	-0.46 ± 0.12	13.4	-0.49 ± 0.13	15.2	-0.45 ± 0.13	12.9
4	8.29 ± 1.33	38.6	7.13 ± 1.34	28.6	7.24 ± 1.33	29.6
5	$(-5.54 \pm 1.53) \times 10^{-3}$	13.1	$(-6.63 \pm 1.53) \times 10^{-3}$	18.7	$(-7.85 \pm 1.57) \times 10^{-3}$	25.0
6	-825 ± 86	92.4	-867 ± 85	103	-883 ± 85	107
$R_{max} = 10000$ km/s (1459 galaxies)						
1	16.3 ± 1.4	143	17.3 ± 1.4	157	17.2 ± 1.4	156
2	1.38 ± 0.18	59.3	1.44 ± 0.18	62.1	1.55 ± 0.18	70.6
3	-0.24 ± 0.11	4.55	-0.26 ± 0.11	5.32	-0.27 ± 0.11	5.54
4	7.62 ± 1.17	42.4	6.80 ± 1.18	33.0	6.99 ± 1.18	35.3
5	$(-5.96 \pm 1.05) \times 10^{-3}$	32.0	$(-6.41 \pm 1.06) \times 10^{-3}$	36.4	$(-6.71 \pm 1.13) \times 10^{-3}$	34.9
6	-785 ± 87	81.8	-817 ± 87	88.2	-845 ± 87	94.9

Table 2 Parameters of the dipole component

Model	R_{max}	D_z	D_x	D_y	D	l , deg	b , deg	σ , km/s
	km/s							
D	8000	147 ± 47	176 ± 58	-136 ± 56	267 ± 65	322	33	1045
DQ	8000	81 ± 50	174 ± 60	-134 ± 58	234 ± 52	323	20	1033
DQO	8000	-59 ± 79	122 ± 104	-103 ± 100	170 ± 95	320	-20	1020
D	10000	139 ± 49	229 ± 59	-106 ± 59	288 ± 64	335	29	1159
DQ	10000	70 ± 53	200 ± 62	-116 ± 62	242 ± 56	330	17	1151
DQO	10000	61 ± 77	85 ± 98	-182 ± 97	210 ± 86	295	17	1138

on the Fig 3 we show the results of other authors: 1 – (Lynden-Bell et al. 1988), 2 – (Hudson et al. 1995), 3 – (Lauer & Postman 1994), 4 – (Parnovsky et al. 2001), 5 – (Dekel et al. 1999), 6 – (da Costa et al. 2000b), 7 – (Hudson et al. 2004), 8 – (Dale et al. 1999), 9 – (Kudrya et al. 2003), 10 – (Watkins et al. 2009). One can see that the result of Parnovsky & Tugay (2004) matches other results better. The new result became closer to the result of Lauer & Postman (1994), but our norm is 2 times less. What caused such a deviation? We found the apex for ‘old’ galaxies used in previous samples but with corrected data for the HI linewidth. At this stage the apex has notably deviated from its original position. This position is shown on Fig. 3 by a square. After adding new data this deviation increased.

Let us discuss how much this deviation affects the global picture of the large-scale motion. Some authors believe that the norm and the apex of the dipole component are the most prominent characteristics of large-scale motion. This was true for very old samples with small depth. Our sample, however, has so large depth that it includes most nearest superclusters like the Great Attractor and Perseus-Pisces. For this

reason, the first few components of the multipole decomposition (3) can not adequately describe all the radial velocity field of the large-scale motion. Moreso, this is hard to do with a single component (4). This phenomenon is due to the fact that a multipole decomposition is capable of describing very-large-scale motion with characteristic scale much larger than the scale of the sample, but can not describe the not-so-large-scale motions with characteristic scale less than the scale of the sample caused by attraction to these superclusters for any reasonable number of components.

As a result, the variation of the subsample depth R_{max} can lead to significant deviation of the vector $\vec{D}$, especially when the subsample boundary crosses a large attractor. At smaller R_{max} values we observe only the increase of radial velocity in the direction of the attractor. At larger R_{max} we see also the backfall. In the framework of DQO-model these effects can be taken into account, but it is too much for the simple D-model to handle. On Fig. 4 we show the variation of the apex position in dependence from the sample depth. One can see that at small R_{max} the apex crosses the galactic plane and approaches the results of most authors

Table 3 Coefficients of the quadrupole component

i	$R_{max} = 8000 \text{ km/s}$				$R_{max} = 10000 \text{ km/s}$			
	DQ-model		DQO-model		DQ-model		DQO-model	
	$q_i, \%$	F_i	$q_i, \%$	F_i	$q_i, \%$	F_i	$q_i, \%$	F_i
1	8.1 ± 1.5	28.8	7.7 ± 1.6	22.6	5.7 ± 1.4	16.6	7.6 ± 1.6	22.6
2	-2.1 ± 1.6	1.6	-1.9 ± 1.7	1.2	-1.9 ± 1.5	1.7	-2.5 ± 1.6	2.3
3	-0.8 ± 2.2	0.1	-1.1 ± 2.3	0.2	2.1 ± 1.9	1.2	-1.2 ± 2.1	0.3
4	0.9 ± 2.3	0.1	1.2 ± 2.5	0.2	4.1 ± 2.1	4.0	2.6 ± 2.5	1.1
5	1.7 ± 2.7	0.4	0.1 ± 2.8	0.0	1.7 ± 2.5	0.5	-0.2 ± 2.7	0.0
Total		6.44		5.10		4.63		4.85

Table 4 Eigenvalues and eigenvectors of the quadrupole tensor

$R_{max},$ km/s	N	Maximum			Minimum			Third axis		
		$Q_1, \%$	l	b	$Q_2, \%$	l	b	$Q_3, \%$	l	b
DQ-model										
3000	292	19.2 ± 3.7	151	−7	$−19.8 \pm 4.2$	80	68	0.6 ± 5.6	58	−20
6000	916	5.6 ± 1.6	186	74	$−8.5 \pm 1.8$	59	10	2.9 ± 2.4	327	13
7000	1068	7.3 ± 1.5	198	81	$−5.9 \pm 1.7$	77	4	$−1.3 \pm 2.3$	347	7
8000	1240	8.1 ± 1.4	142	87	$−6.2 \pm 1.5$	102	−2	$−1.9 \pm 2.1$	12	2
8500	1322	6.8 ± 1.3	68	85	$−5.2 \pm 1.5$	106	−4	$−1.6 \pm 2.0$	16	−3
9000	1379	7.0 ± 1.2	53	79	$−5.6 \pm 1.4$	109	−6	$−1.4 \pm 1.8$	18	−9
9500	1417	6.9 ± 1.2	41	78	$−5.4 \pm 1.4$	109	−4	$−1.5 \pm 1.8$	18	−11
10000	1459	6.3 ± 1.2	55	75	$−4.4 \pm 1.3$	105	−10	$−1.9 \pm 1.8$	13	−11
DQO-model										
3000	292	22.3 ± 4.3	153	−9	$−19.9 \pm 4.3$	80	63	$−2.4 \pm 6.1$	59	−26
6000	916	5.1 ± 1.9	156	71	$−7.8 \pm 1.9$	56	3	2.8 ± 2.7	325	19
7000	1068	6.6 ± 1.8	124	81	$−5.4 \pm 1.7$	75	−6	$−1.2 \pm 2.5$	346	7
8000	1240	7.8 ± 1.6	141	86	$−5.9 \pm 1.6$	90	−3	$−1.9 \pm 2.3$	360	3
8500	1322	7.5 ± 1.6	133	84	$−4.9 \pm 1.5$	92	−5	$−2.6 \pm 2.2$	2	4
9000	1379	8.2 ± 1.6	101	84	$−5.1 \pm 1.5$	90	−6	$−3.1 \pm 2.2$	359	1
9500	1417	7.8 ± 1.6	93	86	$−4.8 \pm 1.5$	99	−4	$−3.0 \pm 2.2$	9	−1
10000	1459	7.8 ± 1.6	120	83	$−5.3 \pm 1.5$	89	−6	$−2.5 \pm 2.2$	359	3

depicted on Fig. 3. An additional argument in favour of the apex deviation being not much of a problem is the following consideration. If in the framework of D-model with the same R_{max} for each galaxy we introduce statistical weights corresponding to the variance of the radial velocity as a function of distance, the apex will correspond to the position $l = 327 \text{ deg}$; $b = 18 \text{ deg}$, which is closer to the result of Parnovsky & Tugay (2004). It is obvious that in such an approach the input from the nearest galaxies will dominate over the more distant galaxies.

Another possible explanation is that the velocity field is not well described by a pure dipole, and the dipole one recovers is unstable with respect to the limiting distance of the sample used. Furthermore, although the dipole is (in principle) a well-defined property of the field, and its mean convergence with distance is predictable within a given cosmological model, the prob-

lem here (as noted previously) is that the multipole basis is not orthogonal for this sample, so that in practice the dipole is not well-defined (in addition to varying stochastically with sample volume). As (Watkins et al. 2009) have shown, this sample bias in the measurement of the dipole is substantial, and has to be taken into account when comparing results from different samples.

As we mentioned before, the usage of the more complex DQO-model allows to describe such phenomena as the backfall. Besides, as it was shown by Parnovsky & Parnowski (2008), the mean deviation of the calculated apex from its true value, caused by measurement errors and the deviations from the Tully-Fisher relationship, is less in the DQO-model in comparison to the D-model. On Fig. 3 the relevant apex position for the DQO-model is denoted by a diamond. We can see that it fairly corresponds to the results of other authors.




Fig. 4 Dependence of bulk motion apex on the sample depth

As it was mentioned above, we used the generalised Tully-Fisher relationship in the form 10. However, we also calculated the bulk motion parameters in the D-model using the classical form 11. We obtained that the bulk motion velocity is 274 km s^{-1} and the apex coordinates are $l = 317$, $b = 36$ for $R_{max} = 10000 \text{ km s}^{-1}$. For $R_{max} = 8000 \text{ km s}^{-1}$ the corresponding values are 254 km s^{-1} , $l = 306$, $b = 41$. As one can see from Table 2 and Fig. 3 these values fall in 1σ confidence areas.

Let us consider the quadrupole component. In Table 3 we present the corresponding coefficients. The traceless quadrupole tensor is described by 5 coefficients:

$$V_r^{qua} = RQ_{ij}n_in_j = R[q_1(n_1^2 - n_3^2) + q_2(n_2^2 - n_3^2) + q_3n_1n_2 + q_4n_1n_3 + q_5n_2n_3]. \quad (13)$$

In Table 3 we also present the errors of these coefficients and their significances according to F-test. The quadrupole coefficients q_i are close to the coefficients obtained by Parnovsky & Tugay (2004). One can see that the significance of all components except the first one is low. Nevertheless, the significance of the first component is high enough for the total significance of the whole quadrupole component to be sufficient. The F values of the quadrupole component are the following: 4.63 for the DQ-model and $R_{max} = 10000 \text{ km s}^{-1}$, 4.85 for the DQO-model and $R_{max} = 10000 \text{ km s}^{-1}$, 6.44 for the DQ-model and $R_{max} = 8000 \text{ km s}^{-1}$, and 5.10 for the DQO-model and $R_{max} = 8000 \text{ km s}^{-1}$. This should be compared to 3.35, which corresponds

Table 5 Coefficients of the octopole component

i	$R_{max} = 8000 \text{ km/s}$		$R_{max} = 10000 \text{ km/s}$	
	$d_i \cdot 10^6$	F_i	$d_i \cdot 10^6$	F_i
1	7.2 ± 3.9	3.5	-4.1 ± 3.0	1.8
2	-2.1 ± 5.2	0.2	-0.4 ± 3.7	0.0
3	-3.7 ± 5.4	0.5	-1.6 ± 4.6	0.1
4	20.3 ± 7.1	8.2	10.6 ± 4.9	4.7
5	-7.8 ± 7.4	1.1	4.3 ± 6.1	0.5
6	24.0 ± 7.6	9.8	23.0 ± 5.5	17.5
7	-13.0 ± 9.6	1.8	-12.5 ± 8.0	2.4
8	10.9 ± 7.8	2.0	15.8 ± 6.1	6.8
9	-17.9 ± 9.0	3.9	-6.2 ± 6.9	0.8
10	33.8 ± 10.2	11.1	25.9 ± 7.6	11.7
Total		4.16		4.37

to the confidence level of 99.5 per cent for 5 degrees of freedom. Thus the quadrupole component is statistically significant according to F-test at 99.5 per cent confidence level both in DQ and DQO-models.

What is the physical sense of the quadrupole component? As one can see from the paper (Parnovsky et al. 2001), it can be naturally combined with the Hubble constant. As a result, we obtain the effective ‘Hubble constant’ depending on direction

$$H(l, b) = H(1 + Q_{ik}n_in_k). \quad (14)$$

Naturally, this effective ‘Hubble constant’ is caused by large-scale collective motion on the sample scale. To estimate the value of its anisotropy we found the eigenvalues and eigenvectors of tensor Q . The three eigenvectors are orthogonal and the sum of three eigenvalues is equal to zero because $\mathbf{Q}$ is a traceless tensor.

In Table 4 we present the maximum, minimum and medium eigenvalues and the corresponding eigenvectors for DQ and DQO-models and different R_{max} . To obtain the eigenvalues of this ‘Hubble constant’ we should add 100 per cent to each of the listed values. For example, for the subsample with $R_{max} = 10000 \text{ km s}^{-1}$ in the framework of DQ-model the maximal ‘Hubble constant’ is equal to 106.3 per cent of the actual Hubble constant, and the minimal – 95.6 per cent. We can see that the anisotropy is weak but statistically significant. The anisotropy is large only for very small scales less than $60h^{-1} \text{ Mpc}$. As one can see from the Table 4, the maximum axis is dominant over the other two. However, the direction of the minimum axis is more stable for both DQ- and DQO-model. In the DQO-model the maximum axis is also stable at $R_{max} > 6000 \text{ km s}^{-1}$. It lies almost in the supergalactic plane. The minimum direction deviates from this plane by approximately 30 degrees of arc.




Fig. 5 Distribution of radial velocity of collective non-Hubble motion at $20h^{-1} \text{ Mpc}$

In the DQ-model the situation is different. As one can see from the Table 4, the variation of the maximum eigenvector is more evident. With the increase of R_{max} the eigenvector starts deviating from the supergalactic plane and at $R_{max} = 10000 \text{ km s}^{-1}$ this deviation reaches approximately 20 degrees of arc. This effect can be explained by the fact that the Local Supercluster in this case constitutes the smaller fraction of the sample.

We also calculated the statistical significance of the octopole component in the DQO-model. The corresponding F values are equal to 4.37 for $R_{max} = 10000 \text{ km s}^{-1}$ and 4.16 for $R_{max} = 8000 \text{ km s}^{-1}$. These values should be compared to the value 2.52, which corresponds to the confidence level of 99.5 per cent for 10 degrees of freedom. Thus, the octopole component is also statistically significant according to F-test at 99.5 per cent confidence level. The same qualitative conclusion was achieved in the paper (Parnovsky & Parnowski 2008). We can also calculate the statistical significance of $\vec{P}$ in eq. (7) as in the paper (Parnovsky et al. 2001; Parnovsky & Tugay 2004). Its F values are equal to 0.21 for $R_{max} = 10000 \text{ km s}^{-1}$ and 2.50 for $R_{max} = 8000 \text{ km s}^{-1}$. This should be compared to 3.78, 2.60, 2.14 and 0.79, which correspond respectively to 99, 95, 90 and 50 per cent confidence levels for 3 degrees of freedom. Thus, for $R_{max} = 10000 \text{ km s}^{-1}$ the octopole trace is insignificant at 50 per cent confidence level, and for $R_{max} = 8000 \text{ km s}^{-1}$ it is significant at 90 per cent confidence level, but this is insufficient to claim that this value is significant for all subsamples.

In Table 5 we present the coefficients of the octopole component, their errors and significances according to F-test. The octopole tensor including the trace $\vec{P}$ is described by 10 coefficients:

$$V_r^{oct} = R^2 \hat{O}_{ijk} n_i n_j n_k = R^2 (d_1 n_1^3 + d_2 n_2^3 + d_3 n_3^3 + d_4 n_1 n_2^2 + d_5 n_1 n_3^2 + d_6 n_2 n_1^2 + d_7 n_2 n_3^2 + d_8 n_3 n_1^2 + d_9 n_3 n_2^2 + d_{10} n_1 n_2 n_3). \quad (15)$$

The coefficients of the octopole component differ from those given by Parnovsky & Tugay (2004) much stronger than the quadrupole ones. Nevertheless, their 1σ confidence areas overlap.

The knowledge of the coefficients of dipole, quadrupole and octopole components allows us to calculate the distribution of the radial velocity of the collective motion of galaxies. On Fig. 5 we present such a distribution for the DQO-model with $R_{max} = 10000 \text{ km s}^{-1}$ and $R = 2000 \text{ km s}^{-1}$, and on Fig. 6 – for $R_{max} = 10000 \text{ km s}^{-1}$ and $R = 8000 \text{ km s}^{-1}$. Note that the amplitudes of the dipole, quadrupole and octopole components are close to each other at $R \sim 6000 \text{ km s}^{-1}$. Thus, for larger distance like in Fig. 6 the octopole component prevails in the velocity field. This figure is characterised by a deep minimum of radial velocity in the direction opposite to the Great Attractor and Shapley concentration. In the direction to these superclusters there is also a minimum caused by a backfall to the Great Attractor. For smaller distances R like in Fig. 5 this minimum becomes a maximum, but the minimum in the opposite direction is deeper than this




Fig. 6 Distribution of radial velocity of collective non-Hubble motion at $80h^{-1} \text{ Mpc}$

maximum. It is clear that this picture is a simplified representation of the velocity field, but still it is much more complex than the simple bulk motion given by the D-model.

5 Estimation of some cosmological parameters and their combinations

As a result, we obtain not only the parameters of the multipole models of the collective motion, but also an estimation of the distance to galaxies using the generalised Tully-Fisher relationship. This allows us to compile a list of peculiar velocities for 1623 galaxies. It can be used to estimate the cosmological parameters Ω_m and σ_8 . For the previous version of the sample it was done in the paper (Parnovsky et al. 2006). We use a method similar to that used by Feldman et al. (2003) but with some changes. We calculate a 2-point correlation function for peculiar velocities, which is approximated by a formula obtained by Juszkiewicz et al. (1999). The formula has a nonlinear dependency on Ω_m and σ_8 , thus these parameters can be estimated by the maximum likelihood method using the correlation function for RFGC galaxies. This yields an estimation of cosmological parameters Ω_m and σ_8 depicted on Fig. 7. There are 2 versions of the approximation formula: a simple one and a complex one. For the complex formula we obtain the values $\Omega_m = 0.26$ and $\sigma_8 = 0.96$. This point is surrounded by 1σ and 2σ confidence areas. They have the banana-like shapes – very long and

narrow curved strips. Judging from the boundaries of these areas, the errors of the cosmological parameters are very large. For this reason, the good agreement in the value of Ω_m between our estimation and other estimations can be just incidental. The simple model yields similar bananas.

Let us compare the obtained confidence areas with more accurate estimations. Let us start from the 3-year WMAP results. On Fig. 8 we reproduce Fig. 7 from the paper (Spergel et al. 2007) with an overlaid plot of 1σ and 2σ confidence areas for the complex model. The blue colour depicts WMAP data, the yellow colour depicts data obtained from weak gravitational lensing, the green colour depicts constraints imposed by both WMAP and lensing, and the red colour depicts our constraints. We can see that our results are similar to the weak lensing data much more than to WMAP data. In any case, these results are obtained with totally different methods, and their good correspondence is very promising.

The results of comparison with 5-year WMAP data are presented on Fig. 9. It is a reproduction of Fig. 19 from the paper (Komatsu et al. 2008) with an overlaid plot of 1σ and 2σ areas for the complex model. It contains constraints set by WMAP, mutual constraints set by WMAP, BAO and supernovae, and our constraints.

The main drawback of our estimation is its graphical form. In many cases it is preferable to deal with numerical constraints. To obtain such constraints we will make use of the long and narrow shape of our confidence areas. If we consider a combination of cosmo-




Fig. 7 1σ and 2σ confidence levels of cosmological constants Ω_m and σ_8 for the complex (solid) and simple (dashed) models

logical parameters of the form $(\Omega_m/0.3)^\alpha \sigma_8$ and α in the range from 0.3 to 0.7 we will obtain an estimation of the combination with better accuracy than either of the two cosmological parameters. The corresponding estimations are given on Fig. 10.

For the previous version of the sample the estimation of such a combination was made in the paper (Parnovsky 2008). Ibid there are many estimations of other authors for different values of α . Of all the values of α the smallest error was reached at $\alpha = 0.35$. On Fig. 11 one can see the boundaries of 1σ confidence areas for the complex and simple models in the $(\Omega_m/0.3)^{0.35} \sigma_8$ against Ω_m coordinates. Both of them fit the constraint $(\Omega_m/0.3)^{0.35} \sigma_8 = 0.91 \pm 0.05$. This result is almost the same as for the previous version of the sample (0.93 ± 0.05). The importance of this α value is caused by the fact that the value $S_8 = (\Omega_m/0.3)^{0.35} \sigma_8$ was used in the paper (Evrard et al. 2008). As the author notes, the low estimation $S_8 = 0.69$ based upon WMAP and SDSS galaxies clusterisation leads to a number of problems. It leads to a contradiction with results of clusters formation modelling. The high estimation $S_8 \sim 0.8 - 0.9$ lifts these problems. The estimation of the share of hot gas following from it corresponds to modern data obtained from the Sunyaev-Zel'dovich effect. Our results support the high estimation. Note that the paper (Evrard et al. 2008) contains many estimations obtained by different authors with different methods. Among them there are both high estimations, close to ours, and low estimations.




Fig. 8 An estimation of cosmological constants Ω_m and σ_8 superimposed on 3-year WMAP results

6 Conclusion

We prepared a new increased and largely revised sample of RFGC galaxies with data about redshifts and HI linewidths. It allowed us to improve the estimations of parameters of the radial velocity field of large-scale collective motions of galaxies using 3 models of its multipole structure. In comparison with the previous versions the main features remained intact, but separate details, for example the apex of the dipole component of the bulk motion in the D-model, significantly changed. As before, the statistical significance according to F-test of both the quadrupole and the octopole components is well above 99.5 per cent. The exact parameters are given in the article for subsamples with maximum distances $100h^{-1} \text{ Mpc}$ and $80h^{-1} \text{ Mpc}$.

The obtained velocity of the bulk motion is in agreement with the Λ CDM model expectation of $\sim 200 \text{ km s}^{-1}$. The value of the bulk motion velocity attracted additional attention after the recent paper of Watkins et al. (2009) who obtained this value as $407 \pm 81 \text{ km s}^{-1}$ at the same scale $100h^{-1} \text{ Mpc}$ as in present article. In this connection some authors immediately started speculating about the challenge to the




Fig. 9 An estimation of cosmological constants Ω_m and σ_8 superimposed on 5-year WMAP results

Λ CDM model and the necessity for inclusion of non-gravitational forces (Ayaita et al. 2009).

For 1623 galaxies we compiled a list of peculiar velocities which will be published in the nearest future. We also intend to use it to determine the distribution of the matter density (including dark matter) at the scales $75h^{-1} \text{ Mpc}$ using POTENT method. This list was also used to estimate the cosmological parameters Ω_m and σ_8 . The obtained constraint is given in graphical form on Fig. 7. The best numerical constraint is given by a combination $S_8 = (\Omega_m/0.3)^{0.35} \sigma_8 = 0.91 \pm 0.05$.

Acknowledgements

We acknowledge the usage of the HyperLeda database (<http://leda.univ-lyon1.fr>).

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

This article is published in the framework of Programme ‘‘Cosmomicrophysics’’ of the National Academy of Sciences of Ukraine.




Fig. 10 Dependence of $(\frac{\Omega_m}{0.3})^\alpha \sigma_8$ on α . The solid line depicts constraint set by both models, the dashed line depicts constraint set only by the complex model only




Fig. 11 1σ confidence areas of S_8 vs. Ω_m dependence. The solid line corresponds to the complex model, the dashed line corresponds to the simple model

References

- Ayaita, Y. et al., 2009 [arXiv:0908.2903]
 Colless, M. et al., 2001, MNRAS, 321, 277
 da Costa, L. N. et al., 2000a, Astron. J., 120, 95
 da Costa, L. N. et al., 2000b, ApJ, 537, L81
 Dale, D. A. et al., 1999, ApJ, 510, L11
 Dekel, A. et al., 1999, ApJ, 522, 1
 Doyle, M. T. et al., 2005, MNRAS, 361, 34
 Evrard, A. E. et al., 2008, ApJ, 672, 122 (astro-ph/0702241)
 Feldman, H. et al., 2003, ApJ, 596, L131 (astro-ph/0305078)
 Haynes, M. P. et al., 1999a, Astron. J., 117, 1668
 Haynes, M. P. et al., 1999b, Astron. J., 117, 2039
 Hudson, D. J., 1964, “Statistics Lectures on Elementary
 Statistics and Probability”, CERN: Geneva
 Hudson, M. J. et al., 1995, MNRAS, 274, 305
 Hudson, M. J. et al., 1999, ApJ, 512, L79
 Hudson, M. J. et al., 2004, MNRAS, 352, 61
 Karachentsev, I. D., 1989, Astron. J., 97, 1566
 Karachentsev, I. D. et al., 1993, Astron. Nachr., 314, 97
 Karachentsev, I. D. et al., 1995, Astron. Nachr., 316, 369
 Karachentsev, I. D. et al., 1999, Bull. SAO, 47, 5 (astro-ph/0305566)■
 Karachentsev, I. D. et al., 2000a, Bull. SAO, 50, 5
 (astro-ph/0107058)
 Karachentsev, I. D. et al., 2000b, Astron. Rep., 44, 150
 Komatsu, E. et al., 2008, ApJS, 180, 330 [arXiv:0803.0547]
 Koribalski, B. S. et al., 2004, Astron. J., 128, 16
 Kudrya, Yu. N. et al., 2003, A&A, 407, 889
 Juskiewicz, R. et al., 1999, ApJ, 518, L25
 Lauer, T. R. & Postman, M., 1994, ApJ, 425, 418
 Lynden-Bell, D. et al., 1988, ApJ, 326, 19
 Mitronova, S. N. et al., 2003, Bull. SAO, 57, 5
 Parnovsky, S. L., 2008, Astron. Lett., 34, 451
 Parnovsky, S. L. & Gaydamaka, O. Z., 2004, Kinematics
 and Physics of Celestial Bodies, 20, 477
 Parnovsky, S. L. & Tugay, A. V., 2004, Astron. Lett., 30,
 357
 Parnovsky, S. L. & Tugay, A. V., 2005, preprint (astro-ph/0510037)■
 Parnovsky, S. L. & Parnowski, A. S., 2008, Astron. Nachr.,
 329, 864
 Parnovsky, S. L. et al., 2001, Astron. Lett., 27, 765
 Parnovsky, S. L. et al., 2006, Ap&SS, 302, 207
 Paturel, G. et al., 2003a, A&A, 412, 45
 Paturel, G. et al., 2003b, A&A, 412, 57
 Sharov, P. Yu. & Parnovsky, S. L., 2006, Astron. Lett., 32,
 287
 Spergel, D. N. et al., 2007, ApJS, 170, 377 (astro-ph/0603449)■
 Springob, C. M. et al., 2005, ApJS, 160, 149
 Springob, C. M. et al., 2007, ApJS, 172, 599 [arXiv:0705.0647]■
 Theureau, G. et al., 2005, A&A, 430, 373
 Tonry, J. L. et al., 1997, ApJ, 475, 399
 Tonry, J. L. et al., 2000, ApJ, 530, 625
 Tully, R. B. & Fisher, J. R., 1977, A&A, 54, 661
 de Vaucouleurs, G. et al., 1991, “Third Reference Catalogue
 of Bright Galaxies”, Springer: Berlin, Heidelberg, New
 York
 Watkins, R. et al., 2009, MNRAS, 392, 743
 Wegner, G. et al., 1996, ApJS, 106, 1